



A screen for sleep and starvation resistance identifies a wake-promoting role for the auxiliary channel *unc79*

Kazuma Murakami ¹, Justin Palermo,¹ Bethany A. Stanhope,¹ Allen G. Gibbs,² and Alex C. Keene ^{1,*}

¹Department of Biological Sciences, Florida Atlantic University, Jupiter, FL 33458, USA and

²Department of Biological Sciences, University of Nevada, Las Vegas, Las Vegas, NV 89154, USA

*Corresponding author: Email: KeeneA@FAU.edu

Abstract

The regulation of sleep and metabolism are highly interconnected, and dysregulation of sleep is linked to metabolic diseases that include obesity, diabetes, and heart disease. Furthermore, both acute and long-term changes in diet potentially impact sleep duration and quality. To identify novel factors that modulate interactions between sleep and metabolic state, we performed a genetic screen for their roles in regulating sleep duration, starvation resistance, and starvation-dependent modulation of sleep. This screen identified a number of genes with potential roles in regulating sleep, metabolism, or both processes. One such gene encodes the auxiliary ion channel UNC79, which was implicated in both the regulation of sleep and starvation resistance. Genetic knockdown or mutation of *unc79* results in flies with increased sleep duration, as well as increased starvation resistance. Previous findings have shown that *unc79* is required in pacemaker for 24-hour circadian rhythms. Here, we find that *unc79* functions in the mushroom body, but not pacemaker neurons, to regulate sleep duration and starvation resistance. Together, these findings reveal spatially localized separable functions of *unc79* in the regulation of circadian behavior, sleep, and metabolic function.

Keywords: sleep; metabolic rate; *Drosophila*; feeding; energy stores; mushroom body

Introduction

Sleep acutely regulates metabolic function, and growing evidence suggests that these processes interact to regulate many biological functions including cognition, physiology, and longevity (Hartmann 1974; Siegel 2005; Joiner 2016; Beckwith and French 2019). At the clinical level, many diseases related to metabolic dysfunction including diabetes, heart disease, and obesity are associated with chronic sleep loss (Taheri et al. 2004; Arble et al. 2015; Reutrakul and Van Cauter 2018). In addition, diet potentially influences sleep duration and quality, indicating that neural systems regulating sleep are sensitive to internal nutrient stores and food availability (Catterson et al. 2010; Grandner et al. 2010, 2014; Linford et al. 2012). Identifying how sleep, diet, and metabolic regulation are interconnected is critical to understanding the fundamental functions of sleep.

Interactions between sleep, feeding, and metabolic regulation are highly conserved between the fruit fly and mammals (Griffith 2013; Yurgel et al. 2015; Beckwith and French 2019; Shafer and Keene 2021). Experimental evolution and artificial selection approaches have revealed a relationship between sleep, feeding, and starvation resistance (Masek et al. 2014; Slocumb et al. 2015; Brown et al. 2019). For example, selection for short-sleeping flies results in reduced energy stores and sensitivity to starvation, while selecting for starvation resistance increases sleep duration (Seugnet et al. 2009; Masek et al. 2014; Slocumb et al. 2015). Furthermore, examining naturally occurring genetic variation in

sleep and starvation resistance in *Drosophila melanogaster* from different geographic localities suggests sleep and starvation resistance are inversely related (Brown et al. 2018; Sarikaya et al. 2020). In addition, shared neural circuits appear to regulate both feeding and starvation resistance, including a role for Insulin-like peptides and the Leucokinin Receptor, and naturally occurring variation in levels of the *foraging* gene impact feeding and starvation resistance (Thimngan et al. 2012; Allen et al. 2017; Zandawala et al. 2018). The interactions between these traits under conditions of experimental evolution raise the possibility that shared genetic factors underlie sleep and starvation resistance.

Energy conservation has long been proposed to be primary function of sleep (Hartmann 1973; Berger and Phillips 1995). *Drosophila* live for only a few days in the absence of food, providing an excellent model to examine the effects of sleep on metabolic regulation and energy conservation (Yurgel et al. 2015; Ly et al. 2018). Quantifying longevity under starvation conditions provides a readout of overall energy stores and metabolic rate (Baldal et al. 2006; Schwasinger-Schmidt et al. 2012). In addition, flies acutely suppress sleep and increase activity in response to starvation, providing a system to investigate acute modulation of sleep and metabolic function (Lee and Park 2004; Keene et al. 2010). Genetic screens and genomic analyses have identified many regulators of sleep, metabolic regulation, and starvation resistance, establishing flies as a model for studying the interactions between these processes (Harbison et al. 2005; Jumbo

Received: February 8, 2021. Accepted: May 25, 2021

© The Author(s) 2021. Published by Oxford University Press on behalf of Genetics Society of America.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Lucioni et al. 2010; Murakami et al. 2016; Sonn et al. 2018). Many of the genes initially identified through screening for short-sleeping mutants have reduced life spans or increased sensitivity to stressors, though the relationship with starvation resistance is less clear (Koh et al. 2008; Bushey et al. 2010; Hill et al. 2018). A complete understanding of how these processes are integrated requires the localization of genes and neurons that regulate sleep and metabolic processes.

The study of sleep in flies has predominantly focused on the role of genes and neurons under fed conditions, leading to the identification of many distinct circuits that promote sleep and wakefulness (Allada and Siegel 2008; Sehgal and Mignot 2011; Ly et al. 2018). There is growing evidence that additional cell types are critical regulators of sleep including multiple classes of glia, endocrine cells, and the fat body (Artiushin et al. 2018; Stahl et al. 2018; Vanderheyden et al. 2018; Yurgel et al. 2018; Ertekin et al. 2020). Furthermore, the genes and neurons regulating sleep can differ based on environmental context (Griffith 2013; Beckwith and French 2019; Shafer and Keene 2021). These studies highlight brain-periphery interactions that are change in response numerous environmental contexts including food availability. Therefore, identifying genetic regulators that impact both sleep and metabolic function requires investigating both neuronal and nonneuronal cell types.

Here, we have performed a genetic screen to identify genetic regulators of sleep and metabolic function, targeting genes ubiquitously to identify factors that function both within the brain and the periphery. Flies were tested in a pipeline that measured sleep parameters under fed and starved conditions, followed by assessment of starvation resistance. This screen identified several candidate genes regulating sleep and metabolic function including the sodium leak channel NALCN accessory subunit *uncoordinated 79* (*unc79*). The gene was first identified in the nematode *Caenorhabditis elegans* in a screen that identified numerous regulators behavior and morphology (Brenner 1974). Both *unc79* and its partner *unc80*, are auxiliary subunits of the drosophila *Narrow Abdomen* (*na*), that is a critical regulator of the circadian clock in *Drosophila* (Lear et al. 2005, 2013; Flourakis et al. 2015). Flies mutant for *na* mutants have altered response to anesthetics in these behavioral assays, deficits in circadian locomotor rhythms, and altered social clustering (Humphrey et al. 2007; Burg et al. 2013). Here, we find additional roles for *unc79* in regulating sleep and metabolic regulation. Furthermore, we provide evidence that *unc79* regulates these processes through independent neural circuit and molecular mechanism from those that govern circadian rhythms.

Methods

Fly husbandry

Flies for behavioral experiments were maintained and tested in humidified incubators at 25°C and under 65% humidity (Powers Scientific). Flies were reared on a 12 hours: 12 hours light-dark cycle for experiments prior to behavioral analysis. All flies were maintained on Nutri-fly *Drosophila* food (Genesee Scientific). All RNAi lines tested were obtained from the Bloomington *Drosophila* Stock Center (Bloomington, IN, USA) (Ni et al. 2008; Perkins et al. 2015) and the Vienna *Drosophila* Resource Center stock (# 45780) was used to validate RNAi phenotypes from the initial screen (Vienna, Austria) (Dietzl et al. 2007) (see Table 1). Bridget Lear (Northwestern) generously provided *unc79*^{F01615} and *unc79*^{F01615} lines (Lear et al. 2013).

Table 1 Circadian phenotypes in *unc79* knockdown flies. Data depict the power and period (length) of rhythms in *unc79* knockdown and associated control lines.

Name	Power	±	Period (hours)	±	N
W ¹¹¹⁸ (+)	145.5	18.38	24.78	0.1326	32
<i>luc</i> ^{RNAi/+}	418	38.99	24.24	0.09269	32
<i>unc79</i> ^{RNAi/+}	207.7	22.14	23.84	0.04861	32
<i>nSyb</i> ^{G4L4>luc} ^{RNAi}	420.4	56.79	25.58	0.02795	32
<i>nSyb</i> ^{G4L4>unc79} ^{RNAi}	45.4	7.049	26.32	0.9583	31
OK107 ^{G4L4>luc} ^{RNAi}	309.8	38.95	23.54	0.09379	32
OK107 ^{G4L4>unc79} ^{RNAi}	129	13.16	23.58	0.1117	30

Behavioral analysis

Drosophila Activity Monitors (DAM; Trikinetics, Waltham, MA, USA) were used for all behavioral analyses. The DAM system detects activity by monitoring infrared beam crossings for each animal (Pfeiffenberger et al. 2010a). These data were used to calculate sleep information by extracting immobility bouts of 5 minutes using the *Drosophila* Sleep Counting Macro (Pfeiffenberger et al. 2010b). All behavioral experiments used 5–7 days old mated female flies unless otherwise noted. We have previously found that females display more robust starvation-induced sleep suppression, and have used mated females in a prior genetic screen (Keene et al. 2010; Murakami et al. 2016). For experiments examining the effects of starvation on sleep, activity was then measured for 24 hours on food, prior to transferring flies into tubes containing 1% agar (Fisher Scientific) at ZT0 and activity was recorded for an additional 48 hours.

To measure starvation resistance, flies were starved on experimental day 2 by transferring them from food 1% agar (Fisher Scientific) individual DAM tubes containing 1% agar. Activity across the first 48 hours on agar was used to measure starvation-induced sleep suppression as previously described (Masek et al. 2014). While previous studies measured starvation-induced sleep suppression on day 1 of starvation, we measured this process for multiple days (Keene et al. 2010). To measure starvation resistance, activity was then recorded until death. Death was manually determined as the last activity time point from the final recorded activity bout for each individual fly.

For experiments quantifying circadian rhythm analysis, locomotor activity under free-running conditions was measured using the DAM system as previously described (Chiu et al. 2010). Individual flies were housed in 10% sucrose DAM tubes instead of standard fly food to prevent larval development that interferes that the circadian assay. Five-day-old adult flies were entrained to light-dark (LD) 12 hours: 12 hours (12:12) cycles for three days, then transferred to constant darkness (DD) for 7–8 days. Locomotor activity data were analyzed using Clocklab software (ActiMetrics, Version 2.72). Individual periods were calculated from 7 to 8 days activity data during DD using chi-square periodogram. Rhythm strength was determined by Fast Fourier Transform (FFT) analysis as previously described (Chiu et al. 2010).

Statistical analysis

Statistical analyses were performed using InStat software (GraphPad Software 6.0). For analysis of sleep, we employed a one- or two-way ANOVA followed by a Tukey's *post hoc* test. For starvation resistance, we applied Kaplan–Meier analysis by grouping each genotype.

Data availability

Fully analyzed data and statistics are available as supplementary files. Supplementary material is available at figshare: <https://www.doi.org/10.25387/g3.14738895>. Raw data will be made available upon request.

Results

We developed a pipeline that measured sleep, starvation-induced sleep suppression, and starvation resistance in individual flies. Sleep was measured for 24 hours on standard food, after which, flies were transferred to agar where they were maintained until death to measure starvation-induced sleep suppression and starvation resistance (Figure 1A). We first validated the pipeline in *w¹¹¹⁸* flies, and found that flies robustly reduce sleep the first 24 hours of starvation and live an average of 48 hours without food (Figure 1C).

To screen for novel regulators of sleep and starvation resistance we ubiquitously knocked down genes by expressing RNAi transgenes from the TRiP Collection under control of the *Actin5C-GAL4* driver (Perkins et al. 2015). Control flies harboring *Actin5C-GAL4* driving *UAS-luciferase-RNAi* (*Act5c>Luc^{RNAi}*), a control with no endogenous targets, flies suppressed sleep during the first day of starvation, and an even greater suppression was observed on day 2 of starvation. To enrich for genes that may be involved in sleep or metabolic function, candidate genes were selected from a genome-wide analysis of polymorphisms and genomic markers of selection in flies selectively bred for starvation resistance (Hardy et al. 2018). Of the 1429 significant genes from this analysis, we identified 914 genes that TRiP RNAi stocks available (Perkins et al. 2015; Hardy et al. 2018). Of these, 299 lines (32.7%) were lethal with ubiquitous knockdown and, therefore, were not screened. In total, we screened 616 lines for sleep, starvation-induced sleep suppression (Figure 1D and Supplementary Table S1). Ubiquitous knockdown of the previously identified Ly-6 transmembrane protein, *qur/sleepless*, resulted in the shortest sleep duration, confirming the ability of the screening procedure to effectively identify genetic regulators of sleep (Koh et al. 2008). To examine the relationship between sleep and starvation resistance we plotted the average for each trait. There was no association between these traits ($r^2 < 0.001$, $P > 0.772$), suggesting sleep and starvation resistance independently regulated (Figure 1D). However, we identified a number of genes where ubiquitous knockdown resulted in increased sleep on food and greater starvation resistance (Figure 1D). We also examined the correlation between genes screened for different sleep parameters. For example, daytime sleep duration is correlated with nighttime sleep duration, suggesting shared genes regulate both processes (Supplementary Figure S1A). We found average bout length was inversely correlated with sleep bout number, suggesting these traits are functionally related (Supplementary Figure S1B). However, no correlation was observed between waking activity and total sleep (Supplementary Figure S1C) suggesting independent regulation of these traits. We chose to focus on the gene encoding for the NALCN auxiliary protein, *unc79* because of the robustness of each phenotype and its role as an essential regulator of circadian rhythms sleep regulation (Lear et al. 2005; Joiner et al. 2013).

To validate the sleep and starvation phenotypes associated with *unc79* we repeated experiments and examined the sleep profile. Flies with ubiquitous knockdown of *unc79* (*Act5c>unc79^{RNAi}*) slept significantly more than control flies (Figure 2, A and B).

Furthermore, while both control groups suppressed sleep during day 1 and 2 of starvation, *Act5c>unc79^{RNAi}* flies did not suppress sleep, suggesting that *unc79* is required for metabolic regulation of sleep (Figure 2, A and B). In addition, starvation resistance was significantly increased in *Act5c>unc79^{RNAi}* flies compared to *Act5c>Luc^{RNAi}* controls (Figure 2C). In agreement with our previous findings, waking activity in female control flies increased starvation, but was unchanged in *Act5C>Unc79^{RNAi}* flies (Supplementary Figure S2A) (Keene et al. 2010). Furthermore, the overall waking activity was elevated in *Act5c>Unc79^{RNAi}* flies compared to *Act5c>Luc^{RNAi}* controls under fed conditions suggesting that the increased sleep in flies deficient for *unc79* is not due to general lethargy.

To confirm that the observed phenotypes are not due to RNAi off targets we tested flies with a genetic mutation in the *unc79* locus. The independent Pbac element insertions in the *unc79* locus slept longer on food and failed to suppress sleep when starved, phenocopying RNAi knockdown (Figure 2, D and E, Lear et al. 2013). Furthermore, *unc79* mutants (*unc79^{F03453}* and *unc79^{F01615}*) survived significantly longer on agar than respective controls (Figure 2F). Analysis of flies heterozygous for the mutation revealed the long sleeping phenotype is semi-dominant (Figure 2, D–F).

We also assessed male flies to determine whether these phenotypes generalize across sexes. Male *unc79* mutants (*unc79^{F03453}* and *unc79^{F01615}*) flies slept longer on food and failed to suppress sleep when starved (Supplementary Figure S2A) and survived longer on agar than respective controls (Supplementary Figure S2B), but the response was attenuated compared to female flies. Waking activity in male flies did not differ between fed and starved groups of *unc79* mutants, while controls increase waking activity (Supplementary Figure S2C). Therefore, ubiquitous RNAi knockdown or genetic mutation of *unc79* results in increased sleep and starvation resistance, and impaired metabolic regulation of sleep.

To localize *unc79* function in metabolism and sleep we first targeted *unc79^{RNAi}* to all neurons using the driver *n-synaptobrevin-GAL4* (*nSyb-GAL4*) (Riabinina et al. 2015). Knockdown in neurons led to flies that slept significantly more than background controls harboring expressing RNAi to luciferase under fed conditions (Figure 3A). Furthermore, flies with pan-neuronal knockdown of *unc79* (*nSyb-GAL4>unc79^{RNAi}*) also did not significantly reduce sleep during starvation and survived significantly longer, suggesting *unc79* functions in neurons to regulate sleep and starvation resistance (Figure 3, A and B). To further localize the function of *unc79* we targeted RNAi to six types of neurons known to modulate sleep. Knockdown in the circadian neurons using *Pdf-GAL4* or *Tim-GAL4* did not affect sleep or starvation resistance, suggesting the effects on sleep and metabolic function are independent of its role in circadian activity (Supplementary Figure S3A). Furthermore, no effect was observed knocking down *unc79* in the sleep promoting central complex (*23E10-GAL4*) or broad classes of peptidergic cells (*C929-GAL4*) (Supplementary Figure S3, A–D). Knockdown of *unc79* selectively in the mushroom bodies (*OK107-GAL4*) increased total sleep, specifically during the day, and resulted in increased starvation resistance (Figure 3, C–F). Therefore, selective knockdown with *OK107-GAL4* largely phenocopies ubiquitous knockdown, raising the possibility that *unc79* functions in the mushroom body to regulate sleep and starvation resistance.

To determine whether the phenotypes observed are specific lobes of the mushroom body, we tested the effects of *unc79* knockdown in the α/β lobes (*c739-GAL4*), α'/β' lobes (*c305a-GAL4*),




Figure 1 Screening for sleep and starvation resistance. (A) Schematic of TRiP Screen. Ubiquitous Act5c GAL4 driver is crossed to TRiP lines, F1 flies were transferred to *Drosophila* Activity Monitor tubes. Sleep was measured for 24 hours on standard food, after which, flies were transferred to agar where they remained until death to quantify starvation-induced sleep suppression and starvation resistance. (B) Control *w¹¹¹⁸* flies suppressed sleep during the first day of starvation, and an even greater suppression was observed on day two of starvation. (C) *w¹¹¹⁸* flies survived approximately 3 days on agar, providing a robust readout sleep and starvation resistance. Total sleep (minutes) is measured over 24 hours period and starvation resistance is measured in hours. (D) Scatter plot for fed total sleep on x-axis plotted (minutes) to starvation resistance on y-axis (hours) [Simple Linear Regression: $F_{(1, 609)} = 0.1265$, $R^2 = 0.0001$, $P > 0.7223$, $N = 616$ fly lines]. Control flies with no endogenous targets Act5c GAL4 drive *luc^{RNAi}* (yellow), lines tested in (gray), *qvr* has the lowest sleep and SR (blue), and *unc79* highest sleep and SR (red).

and the γ lobes (1471 GAL4 drivers, Krashes et al. 2007; Aso et al. 2009). Flies with *unc79* knockdown in α/β lobes and α'/β' lobes fail to suppress sleep compared to *luc* control, while knockdown in the γ lobes increased total sleep duration compared to *luc* control and fail to suppress sleep when starved (Figure 4A). Starvation

resistance is increased in *unc79* knockdown in each mushroom body subtype compared to their respective controls expressing *luciferase-RNAi* (Figure 4B). Therefore, loss of *unc79* function in each subset of mushroom body neurons impacts sleep and metabolic regulation, while selective loss in the γ lobes largely recapitulates




Figure 2 *unc9* RNAi and mutants have increased sleep and starvation resistance. (A) Sleep profiles depicting the average sleep each hour over a 72 hours experiment for *Act5c>luc^{RNAi}* (gray) and *Act5c>unc9^{RNAi}* (red). Flies were on food for day 1, then transferred to agar for days 2 and 3. (B) *Act5c>unc9^{RNAi}* (red) during fed [two-way ANOVA: $F_{(2, 261)} = 8.551, P < 0.0001, N > 39$], starved day 1 [two-way ANOVA: $F_{(2, 261)} = 8.551, P < 0.0001, N > 39$] and starved day 2 [two-way ANOVA: $F_{(2, 261)} = 8.551, P < 0.0001, N > 39$] flies slept significantly longer compared to *Act5c>luc^{RNAi}* (gray) controls. (C) Starvation resistance of *Act5c>unc9^{RNAi}* (red) is significantly higher than *Act5c>luc^{RNAi}* (black) control (Gehan-Breslow-Wilcoxon test: $\chi^2 = 94.42, df = 1, P < 0.0001$). (D) Sleep profile for hourly sleep averages over a 72 hours experiment for *w¹¹¹⁸* (+) (black), *unc9^{F01615}* (red) and *unc9^{F03453}* (maroon) flies are on food for day 1, then transferred to agar for days 2 and 3. (E) Total sleep is greater in *unc9^{F03453}* mutant under fed [maroon, $F_{(2, 651)} = 71.46, P < 0.0001, N \geq 39$], starved day 1 ($P < 0.001; N \geq 39$) and starved day 2 ($P < 0.001; N \geq 39$) conditions compared to control (gray). Total sleep is greater in *unc9^{F01615}* mutant (red) under fed [$F_{(2, 651)} = 71.46, P < 0.0001, N \geq 39$], starved day 1 ($P < 0.001; N \geq 39$), and starved day 2 ($P < 0.001; N \geq 39$) conditions compared to *w¹¹¹⁸* control (gray). (F) Starvation resistance is greater in *unc9^{F03453}* (maroon, Gehan-Breslow-Wilcoxon test: $\chi^2 = 29.1, df = 1, P < 0.0001$) and *unc9^{F01615}* (red, Gehan-Breslow-Wilcoxon test: $\chi^2 = 18.6, df = 1, P < 0.0001$) flies compared to *w¹¹¹⁸* control (gray). All sleep data are violin plots and SR data are survival curves. *** $P < 0.001$, **** $P < 0.0001$.

the full extent of ubiquitous knockdown. These findings reveal that *unc9* is required in all lobes of the mushroom body for proper sleep and metabolic regulation.

To verify that the increase in sleep and starvation resistance is specific to the mushroom bodies, we examined whether

including of the MB-GAL80 transgene reverses the effects of *unc9* knockdown in 1471-GAL4 positive neurons. Expression of GAL80 in the mushroom bodies restores sleep and starvation-induced sleep suppression to control levels to flies with *unc9* knocked down in the γ lobes (Figure 4C). Starvation resistance increased in




Figure 3 Localization of sleep and starvation resistance phenotype to mushroom body. (A) Pan-neuronal knockdown of *unc79* (*nSyb>unc79^{RNAi}*, red) is significantly increased in sleep during fed [two-way ANOVA: $F_{(2, 300)} = 57.82, P < 0.0001, N > 30$], starved day 1 ($P < 0.0001, N > 30$), and starved day 2 ($P < 0.0001, N > 30$); while *nSyb>attp2* (light gray) and *nSyb>luc RNAi* controls shows starvation-induced sleep suppression. (B) Starvation resistance for pan-neuronal knockdown of *unc79* *nSyb>unc79^{RNAi}* (red) is significantly increased compared to *nSyb>attp2* (gray, Gehan-Breslow-Wilcoxon test: $\chi^2 = 42, df = 1, P < 0.0001, N > 30$) and *nSyb>luc RNAi* (light gray, Gehan-Breslow-Wilcoxon test: $\chi^2 = 64.6, df = 1, P < 0.0001, N > 37$) control flies. (C) Sleep profile hourly sleep averages over a 72-hours experiment for mushroom body knockdown of *unc79*. Flies are on food for day 1, then transferred to agar for days 2 and 3. (D) Mushroom body knockdown of *unc79* (*OK107>unc79^{RNAi}*, red) is significantly increased in sleep during fed [two-way ANOVA: $F_{(2, 300)} = 57.82, P < 0.0001, N > 30$] starved day 1 ($P < 0.0001, N > 31$), and starved day 2 ($P < 0.0001, N > 31$); while *Ok107>attp2* (light gray) and *Ok107>luc RNAi* controls shows starvation-induced sleep suppression. (E) Day-time sleep in flies with mushroom body knockdown of *unc79* (*OK107>unc79^{RNAi}*, red) is significantly increased under fed conditions [two-way ANOVA: $F_{(2, 348)} = 43.42, P < 0.0001, N > 30$], starved day 1 ($P < 0.0001, N > 31$), and starved day 2 ($P < 0.0001, N > 31$); while *OK107>attp2* (gray) and *OK107>luc RNAi* (light gray) controls maintain normal daytime sleep. (F) Starvation resistance is increased in *OK107>unc79^{RNAi}* flies compared to *Ok107>attp2* (gray, Gehan-Breslow-Wilcoxon test: $\chi^2 = 47.6, df = 1, P < 0.0001, N > 31$) and *OK107>luc RNAi* (light gray, Gehan-Breslow-Wilcoxon test: $\chi^2 = 50.5, df = 1, P < 0.0001, N > 31$) control flies. All sleep data represent violin plots and SR data are survival curves. * $P < 0.05$; *** $P < 0.001$; **** $P < 0.0001$.

flies with *unc79* selectively knocked down in the (1471-GAL4>*unc79^{RNAi}*). Blocking expression of GAL4 within the mushroom body restored starvation resistance, and partially restored sleep duration on food, confirming that loss of *unc79* in the mushroom body leads to dysregulated sleep (Figure 4D). In addition, the expression of MB-GAL80 restored normal starvation

resistance to 1471-GAL4>*unc79^{RNAi}* flies. These findings validate that the results obtained with *unc79* knockdown using 1471-GAL4 are specific to loss of *unc79* function within the γ -lobe of mushroom bodies.

Previous work has revealed that *unc79* functions within the circadian neurons in association with the *unc80* accessory protein




Figure 4 *unc79* function in the mushroom body γ -lobe to regulate sleep and starvation resistance. (A) Flies with *unc79* knocked down in the $\alpha\beta$ lobes (c739>*unc79*^{RNAi}, pink) fail to suppress starvation during starved day 1 [two-way ANOVA $F_{(10, 1075)} = 28.56, P < 0.0001, N > 53$] and starved day 2 ($P < 0.0001$) compared to control (c739>*luc*^{RNAi}, light gray), while fed day 1 did not differ ($P > 0.1587$). Mushroom body $\alpha'\beta'$ knockdown of *unc79* (c305a>*unc79*^{RNAi}, red) fails to suppress starvation during starved day 1 [two-way ANOVA $F_{(10, 1075)} = 28.56, P < 0.0001, N > 51$] and starved day 2 ($P < 0.0001$) compared to control (c305a>*luc*^{RNAi}, light gray); while fed day 1 did not differ ($P > 0.999$). Mushroom body γ knockdown of *unc79* (1471>*unc79*^{RNAi}, maroon) significantly increase total sleep during fed day [two-way ANOVA $F_{(10, 1075)} = 28.56, P < 0.0001, N > 53$] and fails to suppress sleep on starved day 1 ($P < 0.0001$), and starved day 2 ($P < 0.0001$) compared to control (1471>*luc*^{RNAi}, dark gray). (B) Starvation resistance increased when mushroom body $\alpha\beta$ knockdown of *unc79* (c739>*unc79*^{RNAi}, pink) is significant (Gehan-Breslow-Wilcoxon test: $\chi^2 = 71.36, df = 1, P < 0.0001, N > 63$) compared to control (c739>*luc*^{RNAi}, light gray). Starvation resistance increased when mushroom body $\alpha'\beta'$ knockdown of *unc79* (c305a>*unc79*^{RNAi}, red) is significant (Gehan-Breslow-Wilcoxon test: $\chi^2 = 90.45, df = 1, P < 0.0001, N > 51$) compared to control (c305a>*luc*^{RNAi}, gray). Mushroom body γ knockdown of *unc79* (1471>*unc79*^{RNAi}, maroon) is significant (Gehan-Breslow-Wilcoxon test: $\chi^2 = 124.6, df = 1, P < 0.0001, N > 53$) compared to control (1471>*luc*^{RNAi}, dark gray). (C) Mushroom body GAL80 rescue γ knockdown of *unc79*^{RNAi} (1471>*MB*^{Gal80}; *unc79*^{RNAi}) significantly rescues total sleep compared to γ knockdown of *unc79* (1471>*unc79*^{RNAi}, $P < 0.0432$); while, total sleep for γ knockdown of *unc79* (1471>*unc79*^{RNAi}) is high compares to other control groups (*MB*^{Gal80}; *luc*^{RNAi/+}, $P < 0.0001$; *unc79*^{RNAi/+}, $P < 0.0001$; 1471/+ , $P < 0.0001$; *MB*^{Gal80}; *luc*^{RNAi/+}, $P < 0.0001$; 1471>*MB*^{Gal80}; *luc*^{RNAi}, $P < 0.0001$; and, *MB*^{Gal80}; *unc79*^{RNAi/+}, $P < 0.0001$). Mushroom body GAL80 rescue γ knockdown of *unc79*^{RNAi} (1471>*MB*^{Gal80}; *unc79*^{RNAi}) restored total sleep to controls (*unc79*^{RNAi/+}, $P = 0.3715$; *MB*^{Gal80}; *unc79*^{RNAi/+}, $P = 0.7292$; and 1471>*MB*^{Gal80}; *luc*^{RNAi}, $P = 0.8177$). However, 1471>*MB*^{Gal80}; *unc79*^{RNAi} vs 1471/+ remained significant ($P < 0.0001$). (D) Starvation resistance of mushroom body GAL80 rescue γ knockdown of *unc79*^{RNAi} (1471>*MB*^{Gal80}; *unc79*^{RNAi}) significantly lower (Gehan-Breslow-Wilcoxon test: $\chi^2 = 11.13, df = 1, P < 0.0009, N > 37$) compared to mushroom body γ knockdown of *unc79* (1471>*unc79*^{RNAi}, maroon). All sleep data are violin plots and SR data are survival curves. * $P < 0.05$; *** $P < 0.001$; **** $P < 0.0001$.

and the ion channel *narrow abdomen* (*na*) to maintain locomotor rhythms during constant darkness (Lear et al. 2005, 2013; Moose et al. 2017). To further investigate whether the sleep and circadian phenotypes are controlled by shared or distinct neural circuits, we knocked down additional components of the *unc79* complex in the mushroom bodies and measured the effects on sleep and starvation resistance. We used RNAi lines from the Vienna *Drosophila* Stock Center, for all lines tested. Knockdown of *unc79* in the mushroom bodies phenocopied the effects on sleep and starvation resistance that were observed with the RNAi line used for screening and genetic mutants, confirming these phenotypes.

Knockdown of *na* or *unc80* throughout the mushroom bodies did not increase sleep duration on food or disrupt starvation-induced sleep suppression (Figure 5A). In addition, knockdown of *unc80* and *na* in the mushroom body had little impact on starvation resistance (Figure 5B). These findings raise the possibility that *unc79* functions independently of its canonical complex with *unc80* and *na* to regulate sleep and starvation resistances.

Expression of *unc79* is required within pacemaker neurons to circadian rhythms. Therefore, it is possible that *unc79* functions in distinct populations of neurons to regulate circadian rhythms, from those regulating sleep and metabolic phenotypes. To




Figure 5 MB knockdown of *unc9* RNAi has normal circadian rhythm. (A) Mushroom body knockdown of *unc9* (*OK107>unc9^{RNAi}*) significantly increased sleep [two-way ANOVA: $F(2, 711) = 169, P < 0.0029, N > 40$] compared to control (*unc9^{RNAi/+}*). Mushroom body knockdown of narrow abdomen (*OK107>na^{RNAi}*) did not differ ($P > 0.9999, N > 35$) compared to control (*na^{RNAi/+}*). Mushroom body knockdown of *unc80* (*OK107>unc80^{RNAi}*) did not differ ($P > 0.9964, N > 42$) compared to control (*unc80^{RNAi/+}*). (B) Starvation resistance mushroom body knockdown of *unc9* (*OK107>unc9^{RNAi}*) is increased (Gehan-Breslow-Wilcoxon test: $\chi^2 = 25.95, df = 1, P < 0.0001, N > 40$) compared to control (*unc9^{RNAi/+}*). Starvation resistance mushroom body knockdown of narrow abdomen (*OK107>na^{RNAi}*) is increased (Gehan-Breslow-Wilcoxon test: $\chi^2 = 33.54, df = 1, P < 0.0001, N > 42$) compared to control (*na^{RNAi/+}*). Starvation resistance mushroom body knockdown of narrow abdomen (*OK107>unc80^{RNAi}*) is no different (Gehan-Breslow-Wilcoxon test: $\chi^2 = 2.486, df = 1, P < 0.1148, N > 43$) compared to control (*unc80^{RNAi/+}*). (C-H) Actogram double plot. Female flies were entrained in light-dark cycle for days 1–3 and dark-dark cycles 4–10 days. Panels (A–C, E, and F) have normal rhythm. Circadian rhythm is disrupted when *unc9* is knocked down pan-neuronally, while *unc9* knockdown in mushroom body has restored rhythm. All sleep data are violin plots and SR data are survival curves. ** $P < 0.01$; *** $P < 0.001$.

directly test this possibility, we measured the effects of mushroom body-specific knockdown of *unc79* on free-running activity in entrained animals. As expected, flies with pan-neuronal knockdown are arrhythmic under conditions of constant darkness, while control flies show robust rhythms (Figure 5, C–F). Conversely, knockdown in OK107-expressing cells does not impact circadian activity (Supplementary Figure S4; Figure 5, G and H). Therefore, *unc79* function to regulate circadian rhythms through distinct neural mechanisms that regulate sleep and starvation resistance.

Discussion

Here, we screened by targeting gene function ubiquitously to identify regulators of sleep and metabolic function. Growing evidence suggests sleep is regulated by complex interactions between the brain and periphery, including the findings that mutants impacting fat storage, and communication from the fat body to the brain significantly impact sleep (Thimgan et al. 2010; Slocumb et al. 2015; Ertekin et al. 2020).

We have identified numerous candidate regulators of sleep, including a novel role for *unc79* in the regulation of sleep and metabolic function. *Unc79* and *unc80* are auxiliary subunits of the sodium leak channel *na*, an ortholog of mammalian NALCN family of ion channels (Swayne et al. 2009). A number of functions have been identified for this complex including a role in the regulation of circadian rhythms, and anesthesia sensitivity (Lear et al. 2005; Humphrey et al. 2007). Previous work found that loss of narrow abdomen or *unc79* increased sensitivity to the anesthetics, halothane and isofluorane, and increases sleep (Humphrey et al. 2007; Joiner et al. 2013) consistent with our findings of increased quiescence in *unc79* mutants. Mutation of *unc79* also facilitates the emergence from anesthesia, raising the possibility that loss of *unc79* promotes state transitions, rather than directly impacting isofluorane sensitivity (Joiner et al. 2013). Therefore, suppression of arousal may be involved in anesthesia and sleep (Joiner et al. 2013). Mutation of *na* also impacts a number of complex behaviors including social clustering (the distance maintained between individual flies) (Burg et al. 2013), and light-mediated locomotor activity (Nash et al. 2002). These findings suggest a complex role for *na* and associated *unc79* genes in regulating brain function.

Multiple lines of evidence suggest the role of *unc79* in the regulation of sleep, metabolic regulation of sleep, and starvation resistance is separate from its essential role in regulating circadian rhythms. First, we localize function to the mushroom body, a region that is critical for regulation of sleep and modulation of behavior in accordance with feeding state (Joiner et al. 2006; Pitman et al. 2006; Sitaraman et al. 2015a; Tsao et al. 2018). Selective knockdown of *unc79* in the gamma-lobes alone phenocopies *unc79* mutants. While there is little evidence to date for direct involvement of gamma-lobe neurons in sleep regulation, mushroom body output neurons from the gamma-lobes have been found to regulate sleep (Aso et al. 2014; Sitaraman et al. 2015b). We previously reported that the mushroom bodies are dispensable for starvation-induced sleep suppression, however the manipulations that led to this conclusion involved pharmacological ablation or acute genetic silencing of the mushroom bodies (Keene et al. 2010). Therefore, it is possible that loss of *unc79* function impacts sleep circuitry through a mechanism that would not be detected in flies with the previously applied genetic manipulations. Second, *na* and *unc80*, two components of a complex that interacts with *unc79* to regulate circadian rhythms, are

dispensable for regulation of sleep and starvation resistance in the mushroom bodies. These findings raise the possibility that *unc79* may function independently of its canonical complex with *unc80* and *na*. However, it is important to note that we are unable to validate the efficacy of the RNAi lines tested, and therefore cannot rule out a possible role for *unc79* functioning through *na*. Further work is needed to address how *unc79* functions may modulate mushroom body physiology and sleep circuitry. Studies examining the role of *unc79* in circadian function and anesthesia sensitivity suggests it functions by regulating *na* activity to modulate neural activity (Moose et al. 2017), and it is possible that *unc79* modulates the function of a different ion channel within the mushroom bodies. The structure and function of *na*/NALCN appear to be highly conserved with the channel enhancing neural excitability (Chua et al. 2020). Therefore, imaging activity of neurons within the mushroom bodies of flies lacking *unc79* may be informative.

We identify three independent phenotypes to the mushroom bodies. First, we find knockdown of *unc79* in the mushroom bodies promotes sleep suggesting a wake-promoting role for mushroom bodies. The mushroom bodies contain both wake and sleep-promoting neurons, and genetic ablation or silencing of the mushroom body increases wakefulness (Joiner et al. 2006; Pitman et al. 2006; Sitaraman et al. 2015a). It is possible that loss of *unc79* is functioning in either sleep promoting or wake-promoting neurons to elicit this phenotype. We also identify two independent metabolic phenotypes to the mushroom bodies. In *Drosophila*, the mushroom circuits have been well-defined including the identification of modulatory neurons, and output neurons that modulate sleep (Aso et al. 2014; Haynes et al. 2015; Sitaraman et al. 2015b). Therefore, the identification of *unc79* as a regulator of sleep provides the opportunity to examine how gamma lobe output neurons are regulated by input neurons and impact the physiology of output neurons.

In addition to the sleep phenotypes, we find *unc79* mutants are resistant to starvation. This finding is particularly interesting because the list of genes chosen for the screen derived from those identified in a Genome Wide Analysis Study for factors associated with starvation resistance (Harbison et al. 2004; Hardy et al. 2018). Many different factors contribute to starvation resistance including energy stores, basal metabolic rate, and changes in metabolic rate upon starvation. Animals selected for starvation resistance have elevated sleep and do not suppress sleep when starved (Masek et al. 2014). Therefore, future work studying starvation selected lines, or other populations of outbred fly lines have potential to identify whether variable expression of *unc79* is associated with naturally occurring differences in sleep and metabolic regulation.

We find that *unc79* most potently impacts sleep and starvation resistance in the gamma lobes, suggesting this population is critical for both sleep metabolic regulation. Output neurons from the gamma lobes have been directly implicated in feeding and fat storage supporting the notion that this region is critical for metabolic regulation (Al-Anzi and Zinn 2018). Future work examining the effects of *unc79* deficiency on the physiology and function of mushroom body output neurons may help identify the role of *unc79* in regulating mushroom body circuits that ultimately regulate behavior and metabolic function. Taken together, these findings add to growing evidence that sleep and metabolic function are integrated. The identification of additional genetic factors that regulate the relationship between sleep and nutritional state through behavioral studies will improve our understanding of the strong associations between sleep loss and metabolism-related

diseases. The ubiquitous screen has identified numerous candidate genes that impact sleep, starvation-induced sleep suppression, and starvation resistance, providing candidates that function within and outside of the nervous system. Future study of these genes, such as *unc79*, has potential to advance our understanding of sleep-metabolism interactions and brain-periphery communication.

Acknowledgments

This work was supported by National Institute of Health awards R01HL143790 and R01DC01790 to A.C.K. The authors are grateful to Peter Lewis (FAU) for technical support.

Conflicts of interest

None declared.

Literature cited

- Al-Anzi B, Zinn K. 2018. Identification and characterization of mushroom body neurons that regulate fat storage in *Drosophila*. *Neural Dev.* 13:18.
- Allada R, Siegel JM. 2008. Unearthing the phylogenetic roots of sleep. *Curr Biol.* 18:R670–R679.
- Allen AM, Anreiter I, Neville MC, Sokolowski MB. 2017. Feeding-related traits are affected by dosage of the foraging gene in *Drosophila melanogaster*. *Genetics.* 205:761–773.
- Arble DMDM, Bass J, Behn CDCD, Butler MPMP, Challet E, et al. 2015. Impact of sleep and circadian disruption on energy balance and diabetes: a summary of workshop discussions. *Sleep.* 38:1849–1860.
- Artiushin G, Zhang SL, Tricoire H, Sehgal A. 2018. Endocytosis at the *Drosophila* blood–brain barrier as a function for sleep. *eLife.* 7:e43326.
- Aso Y, Grübel K, Busch S, Friedrich AB, Siwanowicz I, et al. 2009. The mushroom body of adult *Drosophila* characterized by GAL4 drivers. *J Neurogenet.* 23:156–172.
- Aso Y, Sitaraman D, Ichinose T, Kaun KR, Vogt K, et al. 2014. Mushroom body output neurons encode valence and guide memory-based action selection in *Drosophila*. *eLife.* 3:e04580.
- Baldal EA, Brakefield PM, Zwaan BJ. 2006. Multitrait evolution in lines of *Drosophila melanogaster* selected for increased starvation resistance: the role of metabolic rate and implications for the evolution of longevity. *Evolution (N.Y.)*. 60:1435.
- Beckwith EJ, French AS. 2019. Sleep in *Drosophila* and its context. *Front Physiol.* 10:1167.
- Berger RJ, Phillips NH. 1995. Energy conservation and sleep. *Behav Brain Res.* 69:65–73.
- Brenner S. 1974. The genetics of *Caenorhabditis elegans*. *Genetics.* 77:71–94.
- Brown EB, Slocumb ME, Szuperak M, Kerbs A, Gibbs AG, et al. 2019. Starvation resistance is associated with developmentally specified changes in sleep, feeding and metabolic rate. *J Exp Biol.* 222:jeb191049.
- Brown EB, Torres J, Bennick RA, Rozzo V, Kerbs A, et al. 2018. Variation in sleep and metabolic function is associated with latitude and average temperature in *Drosophila melanogaster*. *Ecol Evol.* 8:4084–4097.
- Burg ED, Langan ST, Nash HA. 2013. *Drosophila* social clustering is disrupted by anesthetics and in narrow abdomen ion channel mutants. *Genes Brain Behav.* 12:338–347.
- Bushey D, Hughes KA, Tononi G, Cirelli C. 2010. Sleep, aging, and lifespan in *Drosophila*. *BMC Neurosci.* 11:56.
- Catterson JH, Knowles-Barley S, James K, Heck MMS, Harmar AJ, et al. 2010. Dietary modulation of *Drosophila* sleep-wake behaviour. *PLoS One.* 5:e12062.
- Chiu JC, Low KH, Pike DH, Yildirim E, Edery I. 2010. Assaying locomotor activity to study circadian rhythms and sleep parameters in *Drosophila*. *J Vis Exp.* 34:2157.
- Chua HC, Wulf M, Weidling C, Rasmussen LP, Pless SA. 2020. The NALCN channel complex is voltage sensitive and directly modulated by extracellular calcium. *Sci Adv.* 6:eaaz3154.
- Dietzl G, Chen D, Schnorrer F, Su K-C, Barinova Y, et al. 2007. A genome-wide transgenic RNAi library for conditional gene inactivation in *Drosophila*. *Nature.* 448:151–156.
- Ertekin D, Kirszenblat L, Faville R, van Swinderen B. 2020. Down-regulation of a cytokine secreted from peripheral fat bodies improves visual attention while reducing sleep in *Drosophila*. *PLoS Biol.* 18:e3000548.
- Flourakis M, Kula-Eversole E, Hutchison AL, Han TH, Aranda K, et al. 2015. A conserved bicycle model for circadian clock control of membrane excitability. *Cell.* 162:836–848.
- Grandner MA, Jackson N, Gerstner JR, Knutson KL. 2014. Sleep symptoms associated with intake of specific dietary nutrients. *J Sleep Res.* 23:22–34.
- Grandner MA, Kripke DF, Naidoo N, Langer RD. 2010. Relationships among dietary nutrients and subjective sleep, objective sleep, and napping in women. *Sleep Med.* 11:180–184.
- Griffith LC. 2013. Neuromodulatory control of sleep in *Drosophila melanogaster*: integration of competing and complementary behaviors. *Curr Opin Neurobiol.* 23:819–823.
- Harbison ST, Chang S, Kamdar KP, Mackay TFC. 2005. Quantitative genomics of starvation stress resistance in *Drosophila*. *Genome Biol.* 6:R36.
- Harbison ST, Yamamoto AH, Fanara JJ, Norga KK, Mackay TFC. 2004. Quantitative trait loci affecting starvation resistance in *Drosophila melanogaster*. *Genetics.* 166:1807–1823.
- Hardy CM, Burke MK, Everett LJ, Han MV, Lantz KM, et al. 2018. Genome-wide analysis of starvation-selected *Drosophila melanogaster*-a genetic model of obesity. *Mol Biol Evol.* 35:50–65.
- Hartmann E. 1974. The function of sleep. *Annu Psychoanal.* 2:271–289.
- Hartmann EL. 1973. *The Functions of Sleep*. New Haven: Yale University Press.
- Haynes PR, Christmann BL, Griffith LC. 2015. A single pair of neurons links sleep to memory consolidation in *Drosophila melanogaster*. *eLife.* 4:e03868.
- Hill VM, O'Connor RM, Sissoko GB, Irobunda IS, Leong S, et al. 2018. A bidirectional relationship between sleep and oxidative stress in *Drosophila*. *PLoS Biol.* 16:e2005206.
- Humphrey JA, Hamming KS, Thacker CM, Scott RL, Sedensky MM, et al. 2007. A putative cation channel and its novel regulator: cross-species conservation of effects on general anesthesia. *Curr Biol.* 17:624–629.
- Janett A, Rubin GM, Ngo TTB, Shepherd D, Murphy C, et al. 2012. A GAL4-driver line resource for *Drosophila* neurobiology. *Cell Rep.* 2:991–1001.

- Joiner WJ. 2016. Unraveling the evolutionary determinants of sleep. *Curr Biol.* 26:R1073–R1087.
- Joiner WJ, Crocker A, White BH, Sehgal A. 2006. Sleep in *Drosophila* is regulated by adult mushroom bodies. *Nature.* 441:757–760.
- Joiner WJ, Friedman EB, Hung HT, Koh K, Sowcik M, et al. 2013. Genetic and Anatomical Basis of the Barrier Separating Wakefulness and Anesthetic-Induced Unresponsiveness. *PLoS Genet.* 9:e1003605.
- Jumbo-Lucioni P, Ayroles JF, Chambers MM, Jordan KW, Leips J, et al. 2010. Systems genetics analysis of body weight and energy metabolism traits in *Drosophila melanogaster*. *BMC Genomics.* 11:297.
- Keene ACAC, Duboué ERER, McDonald DMDM, Dus M, Suh GSBGSB, et al. 2010. Clock and cycle limit starvation-induced sleep loss in *drosophila*. *Curr Biol.* 20:1209–1215.
- Koh K, Joiner WJ, Wu MN, Yue Z, Smith CJ, et al. 2008. Identification of SLEEPLESS, a sleep-promoting factor. *Science.* 321:372–376.
- Krashes MJ, Keene AC, Leung B, Armstrong JD, Waddell S. 2007. Sequential use of mushroom body neuron subsets during *drosophila* odor memory processing. *Neuron.* 53:103–115.
- Lear BC, Darrach EJ, Aldrich BT, Gebre S, Scott RL, et al. 2013. UNC79 and UNC80, putative auxiliary subunits of the NARROW ABDOMEN ion channel, are indispensable for robust circadian locomotor rhythms in *Drosophila*. *PLoS One.* 8:e78147.
- Lear BC, Lin JM, Keath JR, McGill JJ, Raman IM, et al. 2005. The ion channel narrow abdomen is critical for neural output of the *Drosophila* circadian pacemaker. *Neuron.* 48:965–976.
- Lee G, Park JH. 2004. Hemolymph sugar homeostasis and starvation-induced hyperactivity affected by genetic manipulations of the adipokinetic hormone-encoding gene in *Drosophila melanogaster*. *Genetics.* 167:311–323.
- Linford NJ, Chan TP, Pletcher SD. 2012. Re-patterning sleep architecture in *Drosophila* through gustatory perception and nutritional quality. *PLoS Genet.* 8:e1002668.
- Ly S, Pack AI, Naidoo N. 2018. The neurobiological basis of sleep: Insights from *Drosophila*. *Neurosci Biobehav Rev.* 87:67–86.
- Masek P, Reynolds L. A, Bollinger WL, Moody C, Mehta A, et al. 2014. Altered regulation of sleep and feeding contribute to starvation resistance in *Drosophila*. *J Exp Biol.* 217:3122–3132.
- Moose DL, Haase SJ, Aldrich BT, Lear BC. 2017. The narrow abdomen ion channel complex is highly stable and persists from development into adult stages to promote behavioral rhythmicity. *Front Cell Neurosci.* 11:159.
- Murakami K, Yurgel MEME, Stahl BABA, Masek P, Mehta A, et al. 2016. Translin is required for metabolic regulation of sleep. *Curr Biol.* 26:972–980.
- Nash HA, Scott RL, Lear BC, Allada R. 2002. An unusual cation channel mediates photic control of locomotion in *Drosophila*. *Curr Biol.* 12:2152–2158.
- Ni JQ, Markstein M, Binari R, Pfeiffer B, Liu LP, et al. 2008. Vector and parameters for targeted transgenic RNA interference in *Drosophila melanogaster*. *Nat Methods.* 5:49–51.
- Perkins LA, Holderbaum L, Tao R, Hu Y, Sopko R, et al. 2015. The transgenic RNAi project at Harvard medical school: resources and validation. *Genetics.* 201:843–852.
- Pfeiffenberger C, Lear BC, Keegan KP, Allada R. 2010a. Locomotor activity level monitoring using the *Drosophila* activity monitoring (DAM) system. *Cold Spring Harb. Protoc.* 5. :pdb.prot5518.
- Pfeiffenberger C, Lear BC, Keegan KP, Allada R. 2010b. Processing sleep data created with the *Drosophila* Activity Monitoring (DAM) System. *Cold Spring Harb Protoc.* 2010:pdb.prot5520.
- Pitman JL, McGill JJ, Keegan KP, Allada R. 2006. A dynamic role for the mushroom bodies in promoting sleep in *Drosophila*. *Nature.* 441:753–756.
- Reutrakul S, Van Cauter E. 2018. Sleep influences on obesity, insulin resistance, and risk of type 2 diabetes. *Metabolism.* 84:56–66.
- Riabinina O, Luginbuhl D, Marr E, Liu S, Wu MN, et al. 2015. Improved and expanded Q-system reagents for genetic manipulations. *Nat Methods.* 12:219–222.
- Sarikaya DP, Cridland J, Tarakji A, Sheehy H, Davis S, et al. 2020. Phenotypic coupling of sleep and starvation resistance evolves in *D. melanogaster*. *BMC Evol Biol.* 20:126.
- Schwasinger-Schmidt TE, Kachman SD, Harshman LG. 2012. Evolution of starvation resistance in *Drosophila melanogaster*: measurement of direct and correlated responses to artificial selection. *J Evol Biol.* 25:378–387.
- Sehgal A, Mignot E. 2011. Genetics of sleep and sleep disorders. *Cell.* 146:194–207.
- Seugnet L, Suzuki Y, Thimngan M, Donlea J, Gimbel SI, et al. 2009. Identifying sleep regulatory genes using a *Drosophila* model of insomnia. *J Neurosci.* 29:7148–7157.
- Shafer OT, Keene AC. 2021. The regulation of *Drosophila* sleep. *Curr Biol.* 31:R38–R49.
- Siegel JM. 2005. Clues to the functions of mammalian sleep. *Nature.* 437:1264–1271.
- Sitaraman D, Aso Y, Jin X, Chen N, Felix M, et al. 2015a. Propagation of homeostatic sleep signals by segregated synaptic microcircuits of the *Drosophila* mushroom body. *Curr Biol.* 25:2517–2527.
- Sitaraman D, Aso Y, Rubin GM, Nitabach MN. 2015b. Control of sleep by dopaminergic inputs to the *Drosophila* mushroom body. *Front Neural Circuits.* 9:73.
- Slocumb MEME, Regalado MJJM, Yoshizawa M, Neely GGG, Masek P, et al. 2015. Enhanced sleep is an evolutionarily adaptive response to starvation stress in *Drosophila*. *PLoS One.* 10:e0131275.
- Sonn JY, Lee J, Sung MK, Ri H, Choi JK, et al. 2018. Serine metabolism in the brain regulates starvation-induced sleep suppression in *Drosophila melanogaster*. *Proc Natl Acad Sci USA.* 115:7129–7134.
- Stahl BA, Peco E, Davla S, Murakami K, Caicedo Moreno NA, et al. 2018. The taurine transporter Eaata2 functions in ensheathing glia to modulate sleep and metabolic rate. *Curr Biol.* 28:3700–3708.
- Swayne LA, Mezghrani A, Varrault A, Chemin J, Bertrand G, et al. 2009. The NALCN ion channel is activated by M3 muscarinic receptors in a pancreatic β -cell line. *EMBO Rep.* 10:873–880.
- Taheri S, Lin L, Austin D, Young T, Mignot E. 2004. Short sleep duration is associated with reduced leptin, elevated ghrelin, and increased body mass index. *PLoS Med.* 1:e62.
- Thimngan MS, Leahy A, Shaw PJ, Donlea J, Sokolowski MB, et al. 2012. Foraging alters resilience/vulnerability to sleep disruption and starvation in *Drosophila*. *Proc Natl Acad Sci USA.* 109:2613–2618.
- Thimngan MS, Suzuki Y, Seugnet L, Gottschalk L, Shaw PJ. 2010. The Perilipin homologue, Lipid storage droplet 2, regulates sleep homeostasis and prevents learning impairments following sleep loss. *PLoS Biol.* 8:e1000466.
- Tsao CH, Chen CC, Lin CH, Yang HY, Lin S. 2018. *Drosophila* mushroom bodies integrate hunger and satiety signals to control innate food-seeking behavior. *eLife.* 7:e35264.

- Vanderheyden WM, Goodman AG, Taylor RH, Frank MG, Van Dongen HPA, et al. 2018. Astrocyte expression of the *Drosophila* TNF-alpha homologue, Eiger, regulates sleep in flies. *PLoS Genet.* 14:e1007724.
- Yurgel ME, Masek P, DiAngelo J, Keene AC. 2015. Genetic dissection of sleep–metabolism interactions in the fruit fly. *J Comp Physiol A Neuroethol Sensory Neural Behav Physiol.* 201:869–877.
- Yurgel ME, Shah KD, Brown EB, Burns C, Bennick RA, et al. 2018. Ade2 functions in the *Drosophila* fat body to promote sleep. *G3 (Bethesda).* 8:3385–3395.
- Zandawala M, Yurgel ME, Liao S, Texada MJ, Rewitz KF, et al. 2018. Modulation of *Drosophila* post-feeding physiology and behavior by the neuropeptide leucokinin. *PLoS Genet.* 14:e1007767.

Communicating editor: M. Arbeitman